

NPI VALUE FTP DIAGNOSIS FILE LAYOUT 2/1/07				
	Length	Begin	END	
Record_type	12	1	12	NSDIAG_TR
COMP	3	13	15	Must equal your component code (085 = ValueOptions)
CASE	10	16	25	NS_ID
BATCH_NO	2	26	27	Filler
TRANS_ENTER_DT	8	28	35	Date transaction was entered (VO system date)
LASTUP_ID	8	36	43	User ID of the person that entered the transaction on your local system (VO probably leave blank)
LOG_TRAN_NO	6	44	49	Use as sort for processing order.
DSM_Flag	1	50	50	Must equal "B"
Filler_A	13	51	63	blank
ID	10	64	73	blank
LASTNM	16	74	89	Consumer last name. Last name must match last name on CARE
LAST_SUF	3	90	92	Suffix to consumer's name, e.g., 'JR', 'SR', 'I', 'II', etc.
FIRSTNM	11	93	103	Consumer first name. First initial must match first initial on CARE.
MIDNM	1	104	104	Consumer middle initial
Filler	9	105	113	Filler
EFF_DT	8	114	121	Must be within the client NS enrollment dates. If a diagnostic data already exists for this consumer with the same decision date, this transaction will be rejected.
FORM_TM	4	122	125	Time recorded on the CARE-DG-1 form. HHMM military format.
REAS_ACT	1	126	126	1=Admission, 2=Reevaluation, 3 = Death, 4 = Discharge.
AXIS_IND	1	127	127	Identifies principle axis (values = 1 or 2)
DSM11	5	128	132	Axis I Level 1. DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 least significant. This field is required if principal Diag Axis equals '1'.
DSM12	5	133	137	Axis I Level 2. DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 least significant.
DSM13	5	138	142	Axis I Level 3. DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 least significant.
DSM14	5	143	147	Axis I Level 4. DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 least Significant.
DSM15	5	148	152	Axis I Level 5. DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 least Significant.
DSM16	5	153	157	Axis I Level 6. DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 least Significant.
DSM21	5	158	162	Axis 2 Level 1. DSM code for personality, specific developmental disorders and mental retardation. Level 1 indicates the most significant diagnosis, level 4 the least significant. This field is required if principal DIAG AXIS equals '2'.

DSM22				Axis 2 Level 2. DSM code for personality, specific developmental disorders and mental retardation. Level 1 indicates the most significant diagnosis, level 4 the least Significant.
	5	163	167	
DSM23				Axis 2 Level 3. DSM code for personality, specific developmental disorders and mental retardation. Level 1 indicates the most significant diagnosis, level 4 the least Significant.
	5	168	172	
DSM24				Axis 2 Level 4. DSM code for personality, specific developmental disorders and mental retardation. Level 1 indicates the most significant diagnosis, level 4 the least significant.
	5	173	177	
ICD31				Axis 3 Level 1. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table conditionally required: If principal diagnosis = MH or reason for action = '3' (DEATH), then Axis III Level I ICD code is required. If no Axis II DX, then use the word 'NONE' to indicate that there is no diagnosis, but that the diagnosis was reviewed at the time of update. Data entry is a required field at update.
	5	178	182	
ICD32				Axis 3 Level 2. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table.
	5	183	187	
ICD33				Axis 3 Level 3. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table.
	5	188	192	
ICD34				Axis 3 Level 4. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table.
	5	193	197	
ICD35				Axis 3 Level 5. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table.
	5	198	202	
ICD36				Axis 3 Level 6. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table.
	5	203	207	
AXIS3_DT				Date of the physician's examination in which the Axis 3 diagnosis was determined. Format = YYYYMMDD. Must be < = decision date.
	8	208	215	
AXIS4				Code for psychosocial stressors. There are up to 9 occurrences (stressors).
	1	216	216	
ABL_CUR	1	217	217	MR field - Blank for Value
ABL_POT	1	218	218	MR field - Blank for Value
AAMD_PRIM	3	219	221	MR field - Blank for Value
AAMD_SEC	3	222	224	MR field - Blank for Value
AAMD_TERT	3	225	227	MR field - Blank for Value
GEN_COMP	2	228	229	MR field - Blank for Value
CRAN_ANOM	2	230	231	MR field - Blank for Value
SENS_IMPAIR	2	232	233	MR field - Blank for Value
PERCEPT	2	234	235	MR field - Blank for Value
CONVULS_DIS	2	236	237	MR field - Blank for Value
PSY_IMPAIR	2	238	239	MR field - Blank for Value

MOTOR_DYS	2	240	241	MR field - Blank for Value
MOTOR_LOC	1	242	242	MR field - Blank for Value
MOTOR_SEV	1	243	243	MR field - Blank for Value
AAMD_DT	8	244	251	MR field - Blank for Value
IQ	3	252	254	MR field - Blank for Value
IQ_TEST	2	255	256	MR field - Blank for Value
IQ_TEST_DT	8	257	264	MR field - Blank for Value
SQ	3	265	267	MR field - Blank for Value
SQ_TEST	2	268	269	MR field - Blank for Value
SQ_TEST_DT	8	270	277	MR field - Blank for Value
AAMD_VERS	2	278	279	MR field - Blank for Value
DSM_VERS	1	280	280	Version of DSM codes used for diagnosis. Must = '4' for DSMIV.
ICD_VERS	1	281	281	Version of ICD codes used for diagnosis. Must = '9'
AXIS5	2	282	283	Indicates a clinician's overall judgment of a person's psychological, social and occupational functioning using the Global Assessment Functioning Scale (GAF Scale) that assesses mental illness. Level of functioning.
AXIS5_PREV	2	284	285	The previous AXIS 5 code. May be blank.
AXIS42	1	286	286	In Axis IV-1 through IV-10, at least 1 value is required for MH diagnosis. See Axis IV decode table. Opt - up to 9 values
AXIS43	1	287	287	DSM IV code
AXIS44	1	288	288	DSM IV code
AXIS45	1	289	289	DSM IV code
AXIS46	1	290	290	DSM IV code
AXIS47	1	291	291	DSM IV code
AXIS48	1	292	292	DSM IV code
AXIS49	1	293	293	DSM IV code
AXIS410	1	294	294	DSM IV code